





PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIMO	Books	
Search Publ	Med 🔽 for		Ž			,	Go CI	ear	
10,046 i.g	Limits	Preview/Index		History	Clipboard	Details	The state of the s		
	Display Al	ostract ▼ Show: 20 ▼ Sort ▼ Ser					d to Text		
	11: Nucle	eic Acids l	Res 1991 N	lov 11:19(21	):5901-5	Re	elated Artic	les. Links	

A thermodynamic study of unusually stable RNA and DNA hairpins.

Antao VP, Lai SY, Tinoco I Jr.

Department of Chemistry, University of California, Berkeley 94720.

About 70% of the RNA tetra-loop sequences identified in ribosomal RNAs from different organisms fall into either (UNCG) or (GNRA) families (where N = A, C. G, or U; and R = A or G). RNA hairpins with these loop sequences form unusually stable tetra-loop structures. We have studied the RNA hairpin GGAC(UUCG)GUCC and several sequence variants to determine the effect of changing the loop sequence and the loop-closing base pair on the thermodynamic stability of (UNCG) tetra-loops. The hairpin GGAG(CUUG)CUCC with the conserved loop G(CUUG)C was also unusually stable. We have determined melting temperatures (Tm), and obtained thermodynamic parameters for DNA hairpins with sequences analogous to stable RNA hairpins with (UNCG). C(GNRA)G, C(GAUA)G, and G(CUUG)C loops. DNA hairpins with (TTCG). (dUdUCG), and related sequences in the loop, unlike their RNA counterparts, did not form unusually stable hairpins. However, DNA hairpins with the consensus loop sequence C(GNRA)G were very stable compared to hairpins with C(TTTT)G or C(AAAA)G loops. The C(GATA)G and G(CTTG)C loops were also extra stable. The relative stabilities of the unusually stable DNA hairpins are similar to those observed for their RNA analogs.

PMID: 1719483 [PubMed - indexed for MEDLINE]

Display Abstract	Show: 20 Sort Send to Text	

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer